



WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

PCT

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(31) International Patent Classification ^a : A61K 31/00		(11) International Publication Number: WO 00/02548 (43) International Publication Date: 20 January 2000 (20.01.00)	
(21) International Application Number: PCT/US99/15533		(74) Agent: MURASHIGE, Kate et al., Morrison & Foerster LLP, 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).	
(22) International Filing Date: 9 July 1999 (09.07.99)		(81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, DE, DK, HU, IL, IN, IS, JP, KP, KR, LC, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SD, SG, SI, SK, TR, TT, US, UZ, VN, AR, IPO patent (CH, OM, KE, LS, MW, SD, SL, SZ, UG, ZW). European patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). OAPI patent (BF, BI, CF, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TO).	
(30) Priority Data: 09/113,947 10 July 1998 (10.07.98) US		Published Without international search report and to be republished upon receipt of that report.	
(63) Related by Continuation (CUN) or Continuation-In-Part (CIP) to Earlier Application US 09/113,947 (CON) 10 July 1998 (10.07.98)		(71) Applicant (for all designated States except US): OSTEO-SCREEN (US/US); Suite 201, 2040 Babcock Road, San Antonio, TX 78229 (US).	
(72) Inventors: and (73) Inventors/Applicants (for US only): MUNDY, Gregory, R. (US/US); 3719 Morgan's Creek, San Antonio, TX 78230 (US); GARRETT, L. Ross (US/US); 16739 Windhammer, San Antonio, TX 78233 (US); RUSSINI, G. (US/US); 5827 N.W. Loop 410 #1224, San Antonio, TX 78238 (US).			
(54) Title: INHIBITORS OF PROTEASOMAL ACTIVITY FOR STIMULATING BONE AND HAIR GROWTH			
(57) Abstract Compounds that inhibit the activity of NF- κ B or inhibit the activity of the proteasome or both promote bone formation and hair growth and are thus useful in treating osteoporosis, bone fracture or deficiency, primary or secondary hyperparathyroidism, periodontal disease or defect, metastatic bone disease, osteolytic bone disease, post-traumatic joint surgery, and post-dental implantation; they also stimulate the production of hair follicles and are thus useful in stimulating hair growth, including hair density, in subject where this is desirable.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Armenia	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Azerbaijan	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GB	Great Britain	LV	Latvia	SE	Sweden
AZ	Azerbaijan	GR	Greece	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	HR	Croatia	MD	Republic of Moldova	TG	Togo
BB	Barbados	CN	China	ME	Montenegro	TM	Turkmenistan
BE	Belgium	CU	Cuba	ML	Republic of Macedonia	TR	Turkey
BF	Burkina Faso	DK	Denmark	MN	Mongolia	TT	Trinidad and Tobago
BG	Bulgaria	EE	Estonia	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BT	Bhutan	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CC	Cocos (Keeling) Islands	JP	Japan	NL	Netherlands	VN	Viet Nam
CF	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Cote d'Ivoire	KR	Republic of Korea	PL	Poland		
CM	Cameroon	KZ	Kazakhstan	PT	Portugal		
CN	China	LA	Laos	RO	Romania		
CU	Cuba	LC	Trinidad and Tobago	RU	Russian Federation		
CZ	Czech Republic	LU	Luxembourg	SD	Sudan		
DE	Germany	LV	Latvia	SE	Sweden		
DK	Denmark	SI	Slovenia	SG	Singapore		
EE	Estonia	SK	Slovakia				

BEST AVAILABLE COPY

INHIBITORS OF PROTEASOMAL ACTIVITY FOR STIMULATING BONE AND HAIR GROWTH

Technical Field

The invention relates to compositions and methods for use in treating skeletal system disorders in a vertebrate at risk for bone loss, and in treating conditions that are characterized by the need for bone growth, in treating fractures, and in treating cartilage disorders. The invention also relates to enhancing hair density and growth. More specifically, the invention concerns the use of inhibitors of proteasomal activity and inhibitors of NF- κ B activity for these purposes.

Background Art

Inhibitors of proteasomal activity, and to some extent inhibitors of NF- κ B activity, have two important physiological effects. First, they are able to enhance bone formation and are thus useful for treating various bone disorders. Second, they stimulate the production of hair follicles and are thus useful in stimulating hair growth, including hair density, in subject where this is desirable.

Effect on Bone

Bone is subject to constant breakdown and resynthesis in a complex process mediated by osteoblasts, which produce new bone, and osteoclasts, which destroy bone. The activities of these cells are regulated by a large number of cytokines and growth factors, many of which have now been identified and cloned.

There is a plethora of conditions which are characterized by the need to enhance bone formation or to inhibit bone resorption. Perhaps the most obvious is the case of bone fractures, where it would be desirable to stimulate bone growth and to hasten and complete bone repair. Agents that enhance bone formation would also be useful in facial reconstruction procedures. Other bone deficit conditions include bone segmental defects, periodontal disease, metastatic bone disease, osteolytic bone disease and conditions where connective tissue repair would be beneficial, such as healing or regeneration of cartilage defects or injury. Also of great significance is the chronic condition of osteoporosis, including age-related osteoporosis and osteoporosis associated with post-menopausal

hormone status. Other conditions characterized by the need for bone growth include primary and secondary hyperparathyroidism, disuse osteoporosis, diabetes-related osteoporosis, and glucocorticoid-related osteoporosis.

There are currently no satisfactory pharmaceutical approaches to managing any of these conditions. Bone fractures are still treated exclusively using casts, braces, anchoring devices and other strictly mechanical means. Further bone deterioration associated with post-menopausal osteoporosis has been treated with estrogens or bisphosphonates, which may have drawbacks for some individuals. Although various approaches have been tried, as further discussed below, there remains a need for additions to the repertoire of agents which can be used to treat these conditions.

Treatment of bone or other skeletal disorders, such as those associated with cartilage, can be achieved either by enhancing bone formation or inhibiting bone resorption or both. A number of approaches have been suggested which relate to bone formation.

Bone tissue is an excellent source for factors which have the capacity for stimulating bone cells. Thus, extracts of bovine bone tissue obtained from slaughterhouses contain not only structural proteins which are responsible for maintaining the structural integrity of bone, but also biologically active bone growth factors which can stimulate bone cells to proliferate. Among these latter factors are transforming growth factor β , the heparin-binding growth factors (e.g., acidic and basic fibroblast growth factor), the insulin-like growth factors (e.g., insulin-like growth factor I and insulin-like growth factor II), and a recently described family of proteins called bone morphogenetic proteins (BMPs). All of these growth factors have effects on other types of cells, as well as on bone cells.

The BMPs are novel factors in the extended transforming growth factor β superfamily. Recombinant BMP2 and BMP4 can induce new bone formation when they are injected locally into the subcutaneous tissues of rats (Vozney *J Molec Reprod Dev* (1992) 32:160-67). These factors are expressed by normal osteoblasts as they differentiate, and have been shown to stimulate osteoblast differentiation and bone nodule formation *in vitro* as well as bone formation *in vivo* (Harris S. *et al. J Bone Miner Res* (1994) 9:855-63). This latter property suggests potential usefulness as therapeutic agents in diseases which result in bone loss.

The cells which are responsible for forming bone are osteoblasts. As osteoblasts differentiate from precursors to mature bone-forming cells, they express and secrete a number of enzymes and structural proteins of the bone matrix, including Type-1 collagen, osteocalcin, osteopontin and alkaline phosphatase. They also synthesize a number of growth regulatory peptides which are stored in the bone matrix, and are presumably responsible for normal bone formation. These growth regulatory peptides include the BMPs (Harris S. *et al.* (1994), *supra*). In studies of primary cultures of fetal rat calvarial osteoblasts, BMPs 1, 2, 3, 4, and 6 are expressed by cultured cells prior to the formation of mineralized bone nodules (Harris S. *et al.* (1994), *supra*). Like alkaline phosphatase, osteocalcin and osteopontin, the BMPs are expressed by cultured osteoblasts as they proliferate and differentiate.

Although the BMPs are potent stimulators of bone formation *in vitro* and *in vivo*, there are disadvantages to their use as therapeutic agents to enhance bone healing. Receptors for the bone morphogenetic proteins have been identified in many tissues, and the BMPs themselves are expressed in a large variety of tissues in specific temporal and spatial patterns. This suggests that BMPs may have effects on many tissues in addition to bone, potentially limiting their usefulness as therapeutic agents when administered systemically. Moreover, since they are peptides, they would have to be administered by injection. These disadvantages impose severe limitations to the development of BMPs as therapeutic agents.

The fluorides, suggested also for this purpose, have a mode of action which may be related to tyrosine phosphorylation of growth factor receptors on osteoblasts, as described, for example, Burgener *et al.* *J Bone Min Res* (1995) 10:164-171, but administration of fluorides is associated with increased bone fragility, presumably due to effects on bone mineralization.

Small molecules which are able to stimulate bone formation have been disclosed in PCT applications WO98/17267 published 30 April 1998, WO97/15308 published 1 May 1997 and WO97/48694 published 24 December 1997. These agents generally comprise two aromatic systems spatially separated by a linker. In addition, PCT application WO98/23460 published 18 June 1998 discloses the use of the class of compounds known as statins in enhancing bone formation. U.S. application Serial No. 09/096,631 filed 12 June 1998 is directed to compounds for stimulating bone growth that

are generally isoprenoid pathway inhibitors. The contents of this application, as well as that of the PCT applications cited above, are incorporated herein by reference.

Other agents appear to operate by preventing the resorption of bone. Thus, U.S. Patent No. 5,280,040 discloses compounds described as useful in the treatment of osteoporosis. These compounds putatively achieve this result by preventing bone resorption.

Wang, G.-J. *et al.*, *J Formos Med Assoc* (1995) 94:589-592 report that certain lipid clearing agents, exemplified by lovastatin and bezafibrate, were able to inhibit the bone resorption resulting from steroid administration in rabbits. There was no effect on bone formation by these two compounds in the absence of steroid treatment. The mechanism of the inhibition in bone resorption observed in the presence of steroids (and the mechanism of the effect of steroid on bone *per se*) is said to be unknown.

An abstract entitled "Lovastatin Prevents Steroid-Induced Adipogenesis and Osteoporosis" by Cui, Q. *et al.* appeared in the Reports of the ASBMR 18th Annual Meeting (September 1996) *J Bone Mineral Res.* (1996) 11(S1):SS10 which reports that lovastatin diminished triglyceride vesicles that accumulated when osteoprogenitor cells cloned from bone marrow stroma of chickens were treated in culture with dexamethasone. Lovastatin was reported to diminish the expression of certain mRNAs and to allow the cells to maintain the osteogenic phenotype after dexamethasone treatment, and chickens that had undergone bone loss in the femoral head as a result of dexamethasone treatment were improved by treatment with lovastatin.

These data are, however, contrary to reports that dexamethasone and other inducers, such as BMPs, induce osteoblastic differentiation and stimulate osteocalcin mRNA (Bellows, C.G., *et al.*, *Develop Biol* (1990) 140:132-38; Rickard, D.J., *et al.*, *Develop Biol* (1994) 161:218-28). In addition, Ducy, P. *et al.*, *Nature* (1996) 382:448-52 have recently reported that osteocalcin deficient mice exhibit a phenotype marked by increased bone formation and bones of improved functional quality, without impairment of bone resorption. Ducy *et al.* state that their data suggest that osteocalcin antagonists may be of therapeutic use in conjunction with estrogen replacement therapy (for prevention or treatment of osteoporosis).

It has also been shown that lovastatin inhibits lipopolysaccharide-induced NF- κ B activation in human mesangial cells. Gujjar, C. *et al.* *Nephrol Dial Transplant* (1996) 11:690-996.

And Its Effect on Hair Growth

Disorders of human hair growth include male pattern baldness, alopecia areata, alopecia induced by cancer chemotherapy and hair thinning associated with aging. These conditions are poorly understood, but nevertheless common and distressing, since hair is an important factor in human social and sexual communication.

Hair follicle regulation and growth are still not well understood, but represent dynamic processes involving proliferation, differentiation and cellular interactions during tissue morphogenesis. It is believed that hair follicles are formed only in early stages of development and not replaced.

Hardy, M.H. *et al. Trans Genet* (1992) 8:55-61 describes evidence that bone morphogenetic proteins (BMPs), members of the TGF β super family, are differentially expressed in hair follicles during development. Harris, S.E. *et al. J Bone Miner Res* (1994) 9:855-863 describes the effects of TGF β on expression of BMP-2 and other substances in bone cells. BMP-2 expression in mature follicles also occurs during maturation and after the period of cell proliferation (Hardy *et al.* (1992, *supra*). As noted, however, by Blessing, M. *et al. Genes and Develop* (1992) 7:204-215, the precise role functional role of BMP-2 in hair follicle maturation remains unclear.

Approaches to treat baldness abound in the U.S. patent literature. See for example U.S. Patent No. 5,767,152 (cyanocarboxylic acid derivatives), U.S. Patent No. 5,824,643 (keratinocyte growth factors) and U.S. Patent No. 5,910,497 (16-pyrazinyl-substitute-4-aza-androstane 5-alpha-reductase isozyme 1 inhibitors). There are many others.

Gat, U. *et al. Cell* (1998) 95:603-614 has demonstrated that β -catenin causes adult epithelial cells to create hair follicles, a surprising result in light of the known inability of mature cells to do so. β -Catenin is known to play a role in cell-cell adhesion and growth factor signal transfection. It is also known that after ubiquitination, β -catenin is degraded by the proteasomes. Orford, K. *et al. J Biol Chem* (1997) 272:24735-24738. At least one gene associated with hair growth (or lack thereof) has also been reported. Ahmed, W. *et al. Science* (1998) 279:720-724.

Two accepted agents currently used for the treatment of hair loss are the antihypertensive drug Minoxidil and the 5 α -reductase inhibitor Finasteride. Neither is entirely satisfactory. Both suffer from modest efficacy and are inconvenient to

administer. A specific, topically active and easy to administer compound with better efficacy than these agents would represent a marked advance.

Proteasomes and NF- κ B

The present invention discloses convenient assays for compounds that will be useful in the treatment of bone disorders and in stimulating hair growth. The assays involve inhibition of the activity of the transcription factor NF- κ B or of the activity of proteasomal proteases, preferably proteasomal proteases. Compounds which inhibit these activities are generally useful in treating bone and hair growth disorders. Compounds that inhibit the production of the transcription factor and these proteases will also be useful in the invention. Their ability to do so can be further confirmed by additional assays.

The proteasome is a noncompartmentalized collection of unrelated proteases which form a common architecture in which proteolytic subunits are self-assembled to form barrel-shaped complexes (for review, see Baumeister *et al.*, *Cell* (1998) 92:367-380. The proteasome contains an array of distinct proteolytic activities inside eukaryotic cells. Compounds which inhibit proteasomal activity also reduce NF- κ B activity by limiting its capacity to be translocated to the nucleus (Barnes, P.J. *et al. New Engl J Med* (1997) 336:1066-1071.

Disclosure of the Invention

The present invention adds to the repertoire of osteogenic and hair growth stimulating agents by providing drugs which would inhibit key proteins and enzymes involved in proteasomal activity and which decrease the activity of the nuclear transcription factor NF- κ B, and thus stimulate bone and hair growth. In accordance with the present invention, we have discovered that inhibition of the functions of the proteasomal proteins and the transcription factor NF- κ B in bone cells leads to increased bone growth and to hair follicle formation and stimulation. Thus, assessing a candidate compound for its ability to inhibit proteasomal proteins or NF- κ B provides a useful means to identify bone and hair growth anabolic agents.

The present specification thus provides methods for identification of osteogenic compounds to stimulate bone growth and compounds that stimulate hair growth by assessing their capacity to inhibit proteasome activity or to inhibit the activity of the transcription factor NF- κ B, preferably to inhibit proteasomal activity. Also useful in the

As employed herein, the term "subject" embraces human as well as other animal species, such as, for example, canine, feline, bovine, porcine, rodent, and the like. It will be understood by the skilled practitioner that the subject is one appropriate to the desirability of stimulating bone growth or hair growth. Thus, in general, for example, stimulation of hair growth will be confined in most instances to animals that would appropriately exhibit such growth.

As used herein, "treat" or "treatment" include a postponement of development of bone deficit symptoms and/or a reduction in the severity of such symptoms that will or are expected to develop. These terms further include ameliorating existing bone or cartilage deficit symptoms, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, preventing or reversing bone resorption and/or encouraging bone growth. Thus, the terms denote that a beneficial result has been conferred on a vertebrate subject with a cartilage, bone or skeletal deficit, or with the potential to develop such deficit.

By "bone deficit" is meant an imbalance in the ratio of bone formation to bone resorption, such that, if unmodified, the subject will exhibit less bone than desirable, or the subject's bones will be less intact and coherent than desired. Bone deficit may also result from fracture, from surgical intervention or from dental or periodontal disease. By "cartilage defect" is meant damaged cartilage, less cartilage than desired, or cartilage that is less intact and coherent than desired. "Bone disorders" includes both bone deficits and cartilage defects.

Representative uses of the compounds identified by the assay of the invention include: repair of bone defects and deficiencies, such as those occurring in closed, open and non-union fractures; prophylactic use in closed and open fracture reduction; promotion of bone healing in plastic surgery; stimulation of bone ingrowth into non-cemented prosthetic joints and dental implants; elevation of peak bone mass in premenopausal women; treatment of growth deficiencies; treatment of periodontal disease and defects, and other tooth repair processes; increase in bone formation during distraction osteogenesis; and treatment of other skeletal disorders, such as age-related osteoporosis, post-menopausal osteoporosis, glucocorticoid-induced osteoporosis or disease osteoporosis and arthritis, or any condition that benefits from stimulation of bone formation. The compounds of the present invention can also be useful in repair of congenital, trauma-induced or surgical resection of bone (for instance, for cancer

methods of the invention are compounds which inhibit the *in situ* production of the enzymes contained in the proteasome or inhibit the production of NF- κ B, preferably of enzymes of the proteasomes. Once a compound found to inhibit these activities has been identified, it can be used in an additional aspect of the invention - a method to stimulate the growth of bone or of hair by contacting suitable cells with the identified compound.

The cellular contact may include *in vivo* administration and the compounds of the invention are thus useful in treating degenerative bone diseases, fractures, dental problems, baldness, alopecia and the like. These methods are performed, according to the present invention, with compounds identified as inhibitors of proteasome activity or inhibitors of the activity of transcription factor NF- κ B, preferably inhibitors of the proteasome enzymes, or inhibitors of the production of the proteasome enzymes or of NF- κ B, preferably of the proteasome enzymes.

Brief Description of the Drawings

Figure 1 shows a diagram of the isoprenoid pathway.

Modes of Carrying Out the Invention

In accordance with the present invention, there are provided methods of treating bone defects (including osteoporosis, fractures, osteolytic lesions and segmental bone defects) in subjects suffering therefrom said method comprising administering to said subject, in an amount sufficient to stimulate bone growth, a compound which inhibits proteasomal activity and function or the activity of the nuclear transcription factor NF- κ B or the production of these proteins.

Also in accordance with the present invention, there are provided methods of treating disorders of hair growth. Disorders of hair growth may be the result of a defect in the ability of existing hair follicles to extrude hair, or may be the result of a deficiency in the number of hair follicles *per se*. "Stimulation of hair growth" refers to increasing the volume of hair in a particular area of a subject whether this is the result of an increased rate of growth in length and/or thickness from the same number of hair follicles, growth proceeding from an enhanced number of hair follicles, or both. The number of hair follicles can be enhanced by further activating existing hair follicles or by stimulating the appearance or proliferation of hair follicles in a particular region of the skin.

treatment), and in cosmetic surgery. Further, the compounds of the present invention can be used for limiting or treating cartilage defects or disorders, and may be useful in wound healing or tissue repair.

5 Conditions which would be benefited by "treating" or "treatment" for stimulation of hair growth include male pattern baldness, alopecia caused by chemotherapy, hair thinning resulting from aging, genetic disorders which result in deficiency of hair coverage, and, in animals, providing additional protection from cold temperatures. Thus, while use in humans may be primarily of cosmetic benefit, use in animals may be therapeutic as well.

10 The compositions of the invention may be administered systemically or locally. For systemic use, the compounds herein are formulated for parenteral (e.g., intravenous, subcutaneous, intramuscular, intraperitoneal, intranasal or transdermal) or enteral (e.g., oral or rectal) delivery according to conventional methods. Intravenous administration can be by a series of injections or by continuous infusion over an extended period.

15 Administration by injection or other routes of discretely spaced administration can be performed at intervals ranging from weekly to once to three times daily. Alternatively, the compounds disclosed herein may be administered in a cyclical manner (administration of disclosed compound; followed by no administration; followed by administration of disclosed compound, and the like). Treatment will continue until the desired outcome is achieved.

20 In general, pharmaceutical formulations will include a compound of the present invention in combination with a pharmaceutically acceptable vehicle, such as saline, buffered saline, 5% dextrose in water, borate-buffered saline containing trace metals or the like. Formulations may further include one or more excipients, preservatives, solubilizers, buffering agents, albumin to prevent protein loss on vial surfaces, lubricants, fillers, stabilizers, etc. Methods of formulation are well known in the art and are disclosed, for example, in Remington's Pharmaceutical Sciences, latest edition, Mack Publishing Co., Easton PA, which is incorporated herein by reference.

25 Pharmaceutical compositions for use within the present invention can be in the form of sterile, non-pyrogenic liquid solutions or suspensions, coated capsules, suppositories, lyophilized powders, transdermal patches or other forms known in the art. Local administration may be by injection at the site of injury or defect, or by insertion or attachment of a solid carrier at the site, or by direct, topical application of a viscous liquid, or the like. For local administration, the delivery vehicle preferably provides a

matrix for the growing bone or cartilage, and more preferably is a vehicle that can be absorbed by the subject without adverse effects.

5 Delivery of compounds herein to wound sites may be enhanced by the use of controlled-release compositions, such as those described in PCT publication WO93/20859, which is incorporated herein by reference. Films of this type are particularly useful as coatings for prosthetic devices and surgical implants. The films may, for example, be wrapped around the outer surfaces of surgical screws, rods, pins, plates and the like. Implantable devices of this type are routinely used in orthopedic surgery. The films can also be used to coat bone filling materials, such as hydroxyapatite blocks, demineralized bone matrix plugs, collagen matrices and the like. In general, a film or device as described herein is applied to the bone at the fracture site. Application is generally by implantation into the bone or attachment to the surface using standard surgical procedures.

10 In addition to the copolymers and carriers noted above, the biodegradable films and matrices may include other active or inert components. Of particular interest are those agents that promote tissue growth or infiltration, such as growth factors. Exemplary growth factors for this purpose include epidermal growth factor (EGF), fibroblast growth factor (FGF), platelet-derived growth factor (PDGF), transforming growth factors (TGFs), parathyroid hormone (PTH), leukemia inhibitory factor (LIF), insulin-like growth factors (IGFs) and the like. Agents that promote bone growth, such as bone morphogenetic proteins (U.S. Patent No. 4,761,471; PCT Publication WO90/11366), osteogenin (Sampath *et al. Proc. Natl. Acad. Sci. USA* (1987) 84:7109-13) and NaF (Tencer *et al. J. Biomed. Mat. Res.* (1989) 23: 571-89) are also preferred. Biodegradable films or matrices include calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyanhydrides, bone or dermal collagen, pure proteins, extracellular matrix components and the like and combinations thereof. Such biodegradable materials may be used in combination with non-biodegradable materials, to provide desired mechanical, cosmetic or tissue or matrix interface properties.

20 Alternative methods for delivery of compounds of the present invention include use of ALZET osmotic minipumps (Alza Corp., Palo Alto, CA); sustained release matrix materials such as those disclosed in Wang *et al.* (PCT Publication WO90/11366); electrically charged dextran beads, as disclosed in Bao *et al.* (PCT Publication WO92/03125); collagen-based delivery systems, for example, as disclosed in Ksander *et*

al. Ann. Surg. (1990) 211(3):288-94; methylcellulose gel systems, as disclosed in Beck *et al. J. Bone Min. Res.* (1991) 6(11):1257-65; alginate-based systems, as disclosed in Edelman *et al. Biomaterials* (1991) 12:619-26 and the like. Other methods well known in the art for sustained local delivery in bone include porous coated metal prostheses that can be impregnated and solid plastic rods with therapeutic compositions incorporated within them.

The compounds of the present invention may also be used in conjunction with agents that inhibit bone resorption. Antiresorptive agents, such as estrogen, bisphosphonates and calcitonin, are preferred for this purpose. More specifically, the compounds disclosed herein may be administered for a period of time (for instance, months to years) sufficient to obtain correction of a bone deficit condition. Once the bone deficit condition has been corrected, the vertebrate can be administered an anti-resorptive compound to maintain the corrected bone condition. Alternatively, the compounds disclosed herein may be administered with an anti-resorptive compound in a cyclical manner (administration of disclosed compound, followed by anti-resorptive, followed by disclosed compound, and the like).

In additional formulations, conventional preparations such as those described below may be used.

Aqueous suspensions may contain the active ingredient in admixture with pharmacologically acceptable excipients, comprising suspending agents, such as methyl cellulose; and wetting agents, such as lecithin, lysolecithin or long-chain fatty alcohols. The said aqueous suspensions may also contain preservatives, coloring agents, flavoring agents, sweetening agents and the like in accordance with industry standards.

Preparations for topical and local application comprise aerosol sprays, lotions, gels and ointments in pharmaceutically appropriate vehicles which may comprise lower aliphatic alcohols, polyglycols such as glycerol, polyethylene glycol, esters of fatty acids, oils and fats, and silicones. The preparations may further comprise antioxidants, such as ascorbic acid or tocopherol, and preservatives, such as p-hydroxybenzoic acid esters.

Parenteral preparations comprise particularly sterile or sterilized products.

Injectable compositions may be provided containing the active compound and any of the well known injectable carriers. These may contain salts for regulating the osmotic pressure.

If desired, the osteogenic agents can be incorporated into liposomes by any of the reported methods of preparing liposomes for use in treating various pathogenic conditions. The present compositions may utilize the compounds noted above incorporated in liposomes in order to direct these compounds to macrophages, monocytes, as well as other cells and tissues and organs which take up the liposomal composition. The liposome-incorporated compounds of the invention can be utilized by parenteral administration, to allow for the efficacious use of lower doses of the compounds. Ligands may also be incorporated to further focus the specificity of the liposomes.

Suitable conventional methods of liposome preparation include, but are not limited to, those disclosed by Bangham, A.D. *et al. J. Mol Biol* (1965) 23:238-252, Olson, F. *et al. Biochim Biophys Acta* (1979) 557:9-23, Szoka, F. *et al. Proc Natl Acad Sci USA* (1978) 75:4194-4198, Kim, S. *et al. Biochim Biophys Acta* (1983) 728:339-348, and Mayer, *et al. Biochim Biophys Acta* (1986) 858:161-168.

The liposomes may be made from the present compounds in combination with any of the conventional synthetic or natural phospholipid liposome materials including phospholipids from natural sources such as egg, plant or animal sources such as phosphatidylcholine, phosphatidylethanolamine, phosphatidylglycerol, sphingomyelin, phosphatidylserine, or phosphatidylinositol and the like. Synthetic phospholipids that may also be used, include, but are not limited to: dimyristoylphosphatidylcholine, dioleoylphosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine, and the corresponding synthetic phosphatidylethanolamines and phosphatidylglycerols. Cholesterol or other sterols, cholesterol hemisuccinate, glycolipids, cerebroside, fatty acids, gangliosides, sphingolipids, 1,2-bis(oleoyloxy)-3-(trimethyl ammonio) propane (DOTAP), N-[1-(2,3-dioleoyl) propyl]-N,N,N-trimethylammonium chloride (DOTMA), and other cationic lipids may be incorporated into the liposomes, as is known to those skilled in the art. The relative amounts of phospholipid and additives used in the liposomes may be varied if desired. The preferred ranges are from about 60 to 90 mole percent of the phospholipid; cholesterol, cholesterol hemisuccinate, fatty acids or cationic lipids may be used in amounts ranging from 0 to 50 mole percent. The amounts of the present compounds incorporated into the lipid layer of liposomes can be varied with the concentration of the lipids ranging from about 0.01 to about 50 mole percent.

The liposomes with the above formulations may be made still more specific for their intended targets with the incorporation of monoclonal antibodies or other ligands specific for a target. For example, monoclonal antibodies to the BMP receptor may be incorporated into the liposome by linkage to phosphatidylethanolamine (PE) incorporated into the liposome by the method of Leserman, L. *et al. Nature* (1980) 288:602-604.

Veterinary uses of the disclosed compounds are also contemplated, as set forth above. Such uses would include treatment of bone or cartilage deficits or defects associated with hair or fur in domestic animals, livestock and thoroughbred horses.

The compounds of the present invention may be used to stimulate growth of bone-forming cells or their precursors, or to induce differentiation of bone-forming cell precursors, either *in vitro* or *ex vivo*. The compounds described herein may also modify a target tissue or organ environment, so as to attract bone-forming cells to an environment in need of such cells. As used herein, the term "precursor cell" refers to a cell that is committed to a differentiation pathway, but that generally does not express markers or function as a mature, fully differentiated cell. As used herein, the term "mesenchymal cells" or "mesenchymal stem cells" refers to pluripotent progenitor cells that are capable of dividing many times, and whose progeny will give rise to skeletal tissues, including cartilage, bone, tendon, ligament, marrow stroma and connective tissue (see A. Caplan *J. Orthop. Res.* (1991) 9:641-50). As used herein, the term "osteogenic cells" includes osteoblasts and osteoblast precursor cells. More particularly, the disclosed compounds are useful for stimulating a cell population containing marrow mesenchymal cells, thereby increasing the number of osteogenic cells in that cell population. In a preferred method, hematopoietic cells are removed from the cell population, either before or after stimulation with the disclosed compounds. Through practice of such methods, osteogenic cells may be expanded. The expanded osteogenic cells can be infused (or reinfused) into a vertebrate subject in need thereof. For instance, a subject's own mesenchymal stem cells can be exposed to compounds of the present invention *ex vivo*, and the resultant osteogenic cells could be infused or directed to a desired site within the subject, where further proliferation and/or differentiation of the osteogenic cells can occur without immunorejection. Alternatively, the cell population exposed to the disclosed compounds may be immortalized human fetal osteoblastic or osteogenic cells. If such cells are infused or implanted in a vertebrate subject, it may be advantageous to "immunoprotect"

these non-self cells, or to immunosuppress (preferably locally) the recipient to enhance transplantation and bone or cartilage repair.

As stated above, the compounds of the present invention may also be used to stimulate the growth of hair either by enhancing its rate of formation from existing follicles, stimulating inactive follicles, effecting the production of additional hair follicles or some combination of the foregoing, or by any other mechanism that may or may not presently be understood.

Within the present invention, an "effective amount" of a composition is that amount which produces a statistically significant effect. For example, an "effective amount" for therapeutic uses is the amount of the composition comprising an active compound herein required to provide a clinically significant increase in healing rates in fracture repair; reversal of bone loss in osteoporosis; reversal of cartilage defects or disorders; prevention or delay of onset of osteoporosis; stimulation and/or augmentation of bone formation in fracture non-unions and distraction osteogenesis; increase and/or acceleration of bone growth into prosthetic devices; and repair of dental defects. An "effective amount" for uses in stimulating hair growth is that amount which provides the desired effect in terms of length or density of hair. Such effective amounts will be determined using routine optimization techniques and are dependent on the particular condition to be treated, the condition of the patient, the route of administration, the formulation, and the judgment of the practitioner and other factors evident to those skilled in the art. The dosage required for the compounds of the invention (for example, in osteoporosis where an increase in bone formation is desired) is manifested as a statistically significant difference in bone mass between treatment and control groups. This difference in bone mass may be seen, for example, as a 5-20% or more increase in bone mass in the treatment group. Other measurements of clinically significant increases in healing may include, for example, tests for breaking strength and tension, breaking strength and torsion, 4-point bending, increased connectivity in bone biopsies and other biomechanical tests well known to those skilled in the art. General guidance for treatment regimens is obtained from experiments carried out in animal models of the disease of interest. Differences between successfully treated subjects and controls with regard to stimulation of hair growth can generally be ascertained by direct observation. The dosage of the compounds of the invention will vary according to the extent and severity of the need for treatment, the activity of the administered compound, the

general health of the subject, and other considerations well known to the skilled artisan. Generally, they can be administered to a typical human on a daily basis as an oral dose of about 0.1 mg/kg-1000 mg/kg, and more preferably from about 1 mg/kg to about 200 mg/kg. The parenteral dose will appropriately be 20-100% of the oral dose. While oral administration may be preferable in most instances where the condition is a bone deficit (for reasons of ease, patient acceptability, and the like), alternative methods of administration may be appropriate for selected compounds and selected defects or diseases. While topical administration is generally preferable for stimulating hair growth, as generally only local effects are desired, systemic treatment may be preferable in some instances as well.

Assays for Compounds Useful in the Invention

Assays for assessing the ability of a compound to inhibit proteasomal activity and for inhibitors of NF- κ B activity are well known in the art. Two typical, but nonlimiting assays are described below.

Assessment of Proteasomal Activity

Proteasomal activity is measured by an increase in cytoplasmic ubiquitinated protein complexes, as assessed by Western blotting using an anti-ubiquitin antibody.

MG-63 cells are grown in confluency in alpha MEM media and 10% fetal calf serum (FCS). Cells are then treated for 24 hours with specific compounds. Following the indicated treatments, cells are scraped with a disposable scraper, washed twice with phosphate saline solution (137 mM NaCl, 10 mM d-glucose, 4 mM KCl, 0.5 mM Na_2HPO_4 , 0.1 mM KH_2PO_4), centrifuged, and the resulting pellet is suspended in the sample buffer containing 2% SDS, pH 6.75. The samples are heated and the concentration of total protein calculated by means of Micro bicinchoninic acid (BCA) Protein Assay Kit (Pierce, Rockford, IL/USA). The samples are diluted to obtain a final protein concentration of 2 mg/ml, supplemented with 10% 2-mercaptoethanol, 1% bromophenol blue and run on a 4-15% SDS-PAGE. Resulting gels are Western blotted with anti-ubiquitin rabbit polyclonal antibody (diluted 1:100; Sigma, St. Louis, MO/USA). The samples are visualized with horse-radish peroxidase coupled anti-rabbit IgG antibodies (Amersham Corp., Arlington Heights, IL/USA) using ECL detection kits (Amersham Corp.).

NF- κ B Activity Assays

Cells are treated with different concentrations of compounds, and nuclear extracts prepared. Briefly, cells are washed with phosphate-buffered saline, and resuspended in lysis buffer (0.6% Nonidet P-40, 150 mM NaCl, 10 mM Tris-HCl, pH 7.9, 1 mM EDTA, 0.5 mM DTT and a cocktail of protease inhibitors (Complete (TM), Boehringer Mannheim). After incubation on ice for 15 min, nuclei are collected by centrifugation. The pellet is resuspended in nuclear extraction buffer (10 mM Hepes, pH 7.9, 420 mM NaCl, 0.1 mM EDTA, 1.5 mM MgCl_2 , 0.5 mM DTT, protease inhibitors (Complete (TM), Boehringer Mannheim), 25% glycerol), and incubated at 4 degrees C for 30 min. The supernatant is collected and dialyzed in a buffer containing 10 mM TrisOHCl, pH 7.5, 50 mM NaCl, 5 mM MgCl_2 , 1 mM EDTA, 1 mM DTT, and 20% glycerol. After dialysis, the nuclear extract is centrifuged to remove precipitated proteins, and aliquots are stored at -70 C. Protein concentration in the nuclear extracts is measured by the method of Bradford using a dye-binding assay kit (Bio-Rad).

The probe for electrophoretic mobility shift assays is a 32P-labeled double-stranded oligonucleotide containing the consensus sequence specific for NF- κ B (Promega). Nuclear extracts (5 ug) are pre-incubated in 20- μ l reaction mixtures containing 10 mM Tris-HCl, pH 7.5, 50 mM NaCl, 2.5 mM DTT, 0.5 mM EDTA, 1 mM MgCl_2 , 4% glycerol, and 5 ug of poly (dl-dC). After 10 min at room temperature, 10-20 fmol of probe is added, and incubated further for 20 min. DNA-protein complexes are separated from free oligonucleotides on a 5% polyacrylamide/0.5X TBE gel (45 mM Tris-HCl, 45 mM boric acid, 1 mM EDTA). After electrophoresis, gels are dried and autoradiographed.

Assays for Production Inhibition

Compounds which inhibit the production of the enzymes having proteasomal activity or of NF- κ B can be assessed by measuring the level of production of these proteins in the presence and absence of candidate compounds. The levels of production can be readily measured in *in vitro* systems using, for example, immunoassays for the level of protein produced. The levels of such proteins can also be assessed by utilizing, for example, methionine labeling and size separation of proteins in the cells to be assessed. In order to effect a convenient level of protein production for measurement, it

is advantageous to use recombinant expression systems for the relevant enzymes or the NF- κ B so that substantial amounts are produced.

Typical approaches to inhibiting the production of NF- κ B or proteasome enzymes include the use of antisense technology or formation of triplexes with double-stranded forms of nucleotide sequences relevant in the expression of the genes. In addition, various small molecules may also inhibit this production.

Screening Assays - Bone

The osteogenic activity of the compounds used in the methods of the invention can be verified using *in vitro* screening techniques, such as the assessment of transcription of a reporter gene coupled to a bone morphogenetic protein-associated promoter or in alternative assays.

ABA Screening Assay

A rapid throughput screening test for compounds that stimulate bone formation by demonstration that they are capable of stimulating expression of a reporter gene linked to a BMP promoter (a surrogate for the production of bone morphogenetic factors that are endogenously produced) is described in U.S. Application Serial No. 08/458,434, filed 2 June 1995, the entire contents of which are incorporated herein by reference. This assay is also described as a portion of a study of immortalized murine osteoblasts (derived from a mouse expressing a transgene composed of a BMP2 promoter driving expression of T-antigen) in Ghosh-Choudhery, N. *et al. Endocrinology* (1996) 137:331-39. In this study, the immortalized cells were stably transfected with a plasmid containing a luciferase reporter gene driven by a mouse BMP2 promoter (-2736/114 bp), and responded in a dose-dependent manner to recombinant human BMP2.

Briefly, the assay utilizes cells transformed permanently or transiently with constructs in which the promoter of a bone morphogenetic protein, specifically BMP2 or BMP4, is coupled to a reporter gene, typically luciferase. These transformed cells are then evaluated for the production of the reporter gene product; compounds that activate the BMP promoter will drive production of the reporter protein, which can be readily assayed. Many thousands of compounds have been subjected to this rapid screening technique, and only a very small percentage are able to elicit a level of expression of reporter gene 5-fold greater than that produced by vehicle. Compounds that activate the

BMP promoter fall into groups, where members of each group share certain structural characteristics not present in inactive compounds. The active compounds ("BMP promoter-active compounds" or "active compounds") are useful in promoting bone or cartilage growth, and thus in the treatment of vertebrates in need of bone or cartilage growth.

BMP promoter-active compounds can be examined in a variety of other assays that test specificity and toxicity. For instance, non-BMP promoters or response elements can be linked to a reporter gene and inserted into an appropriate host cell. Cytotoxicity can be determined by visual or microscopic examination of BMP promoter- and/or non-BMP promoter-reporter gene-containing cells, for instance. Alternatively, nucleic acid and/or protein synthesis by the cells can be monitored. For *in vivo* assays, tissues may be removed and examined visually or microscopically, and optionally examined in conjunction with dyes or stains that facilitate histologic examination. In assessing *in vivo* assay results, it may also be useful to examine biodistribution of the test compound, using conventional medicinal chemistry/animal model techniques.

Neonatal Mouse Calvaria Assay (*In vitro*)

An assay for bone resorption or bone formation is similar to that described by Gowen M. & Mundy G. *J Immunol* (1986) 136:2478-82. Briefly, four days after birth, the front and parietal bones of ICR Swiss white mouse pups are removed by microdissection and split along the sagittal suture. In an assay for resorption, the bones are incubated in BGJb medium (Irvine Scientific, Santa Ana, CA) plus 0.02% (or lower concentration) β -methylcyclodextrin, wherein the medium also contains test or control substances. The medium used when the assay is conducted to assess bone formation is Fitton and Jackson Modified BGJ Medium (Sigma) supplemented with 6 μ g/ml insulin, 6 μ g/ml transferrin, 6 ng/ml selenous acid, calcium and phosphate concentrations of 1.25 and 3.0 mM, respectively, and ascorbic acid to a concentration of 100 μ g/ml is added every two days. The incubation is conducted at 37°C in a humidified atmosphere of 5% CO₂ and 95% air for 96 hours.

Following this, the bones are removed from the incubation media and fixed in 10% buffered formalin for 24-48 hours, decalcified in 14% EDTA for 1 week, processed through graded alcohols; and embedded in paraffin wax. Three μ m sections of the

calvaria are prepared. Representative sections are selected for histomorphometric assessment of bone formation or bone resorption. Bone changes are measured on sections cut 200 µm apart. Osteoblasts and osteoclasts are identified by their distinctive morphology.

Other auxiliary assays can be used as controls to determine non-BMP promoter-mediated effects of test compounds. For example, mitogenic activity can be measured using screening assays featuring a serum-response element (SRE) as a promoter and a luciferase reporter gene. More specifically, these screening assays can detect signaling through SRE-mediated pathways, such as the protein kinase C pathway. For instance, an osteoblast activator SRE-luciferase screen and an insulin mimetic SRE-luciferase screen are useful for this purpose. Similarly, test compound stimulation of cAMP response element (CRE)-mediated pathways can also be assayed. For instance, cells transfected with receptors for PTH and calcitonin (two bone-active agents) can be used in CRE-luciferase screens to detect elevated cAMP levels. Thus, the BMP promoter specificity of a test compound can be examined through use of these types of auxiliary assays.

In vivo Assay of Effects of Compounds on Murine Calvarial Bone Growth

Male ICR Swiss white mice, aged 4-6 weeks and weighing 13-26 gm, are employed, using 4-5 mice per group. The calvarial bone growth assay is performed as described in PCT application WO95/24211, incorporated by reference. Briefly, the test compound or appropriate control vehicle is injected into the subcutaneous tissue over the right calvaria of normal mice. Typically, the control vehicle is the vehicle in which the compound was solubilized, and is PBS containing 5% DMSO or is PBS containing Tween (2 µl/10 ml). The animals are sacrificed on day 14 and bone growth measured by histomorphometry. Bone samples for quantitation are cleaned from adjacent tissues and fixed in 10% buffered formalin for 24-48 hours, decalcified in 14% EDTA for 1-3 weeks, processed through graded alcohols, and embedded in paraffin wax. Three to five µm sections of the calvaria are prepared, and representative sections are selected for histomorphometric assessment of the effects on bone formation and bone resorption. Sections are measured by using a camera lucida attachment to trace directly the microscopic image onto a digitizing plate. Bone changes are measured on sections cut 200 µm apart, over 4 adjacent 1x1 mm fields on both the injected and noninjected sides of

the calvaria. New bone is identified by its characteristic woven structure, and osteoclasts and osteoblasts are identified by their distinctive morphology. Histomorphometry software (OsteoMeasure, Osteometrix, Inc., Atlanta) is used to process digitizer input to determine cell counts and measure areas or perimeters.

Typical treatment regimens for testing utilize application of the compound to be tested over several days of repeated administration.

Additional In Vivo Assays - Bone

Lead compounds can be further tested in intact animals using an *in vivo* dosing assay. Prototypical dosing may be accomplished by subcutaneous, intraperitoneal or oral administration, and may be performed by injection, sustained release or other delivery techniques. The time period for administration of test compound may vary (for instance, 28 days as well as 35 days may be appropriate). An exemplary, *in vivo* oral or subcutaneous dosing assay may be conducted as follows:

In a typical study, 70 three-month-old female Sprague-Dawley rats are weight-matched and divided into seven groups, with ten animals in each group. This includes a baseline control group of animals sacrificed at the initiation of the study; a control group administered vehicle only; a PBS-treated control group; and a positive control group administered a compound (non-protein or protein) known to promote bone growth. Three dosage levels of the compound to be tested are administered to the remaining three groups.

Briefly, test compound, positive control compound, PBS, or vehicle alone is administered subcutaneously once per day for 35 days. All animals are injected with calcitonin nine days and two days before sacrifice (two injections of calcitonin administered each designated day). Weekly body weights are determined. At the end of the 35-day cycle, the animals are weighed and bled by orbital or cardiac puncture. Serum calcium, phosphate, osteocalcin, and CBCs are determined. Both leg bones (femur and tibia) and lumbar vertebrae are removed, cleaned of adhering soft tissue, and stored in 70% ethanol for evaluation, as performed by peripheral quantitative computed tomography (pQCT; Ferretti, *J. Bone* (1995) 17:353S-64S), dual energy X-ray absorptiometry (DEXA; Laval-Jeantet A. *et al. Calcif Tissue Int'l* (1995) 56:14-18; J. Casez *et al. Bone and Mineral* (1994) 26:61-68) and/or histomorphometry. The effect of test compounds on bone remodeling can thus be evaluated.

Lead compounds can also be tested in acute ovariectomized animals (prevention model) using an *in vivo* dosing assay. Such assays may also include an estrogen-treated group as a control. An exemplary subcutaneous dosing assay is performed as follows:

In a typical study, 80 three-month-old female Sprague-Dawley rats are weighed, matched and divided into eight groups, with ten animals in each group. This includes a baseline control group of animals sacrificed at the initiation of the study; three control groups (sham ovariectomized (sham OVX) + vehicle only; ovariectomized (OVX) + vehicle only; PBS-treated OVX); and a control OVX group that is administered a compound known to promote bone growth. Three dosage levels of the compound to be tested are administered to the remaining three groups of OVX animals.

Since ovariectomy (OVX) induces hyperphagia, all OVX animals are pair-fed with sham OVX animals throughout the 35 day study. Briefly, test compound, positive control compound, PBS, or vehicle alone is administered orally or subcutaneously once per day for 35 days. Alternatively, test compound can be formulated in implantable pellets that are implanted for 35 days, or may be administered orally, such as by gastric gavage. All animals, including sham OVX/vehicle and OVX/vehicle groups, are injected intraperitoneally with calcein nine days and two days before sacrifice (two injections of calcein administered each designated day, to ensure proper labeling of newly formed bone). Weekly body weights are determined. At the end of the 35-day cycle, the animals' blood and tissues are processed as described above.

Lead compounds may also be tested in chronic OVX animals (treatment model). An exemplary protocol for treatment of established bone loss in ovariectomized animals that can be used to assess efficacy of anabolic agents may be performed as follows.

Briefly, 80 to 100 six month old female, Sprague-Dawley rats are subjected to sham surgery (sham OVX) or ovariectomy (OVX) at time 0, and 10 rats are sacrificed to serve as baseline controls. Body weights are recorded weekly during the experiment. After approximately 6 weeks (42 days) or more of bone depletion, 10 sham OVX and 10 OVX rats are randomly selected for sacrifice as depletion period controls. Of the remaining animals, 10 sham OVX and 10 OVX rats are used as placebo-treated controls. The remaining OVX animals are treated with 3 to 5 doses of test drug for a period of 5 weeks (35 days). As a positive control, a group of OVX rats can be treated with an agent such as PTH, a known anabolic agent in this model (Kimmel *et al. Endocrinology* (1993) 132:1577-84). To determine effects on bone formation, the following procedure can be

followed. The femurs, tibiae and lumbar vertebrae 1 to 4 are excised and collected. The proximal left and right tibiae are used for pQCT measurements, cancellous bone mineral density (BMD) (gravimetric determination), and histology, while the midshaft of each tibiae is subjected to cortical BMD or histology. The femurs are prepared for pQCT scanning of the midshaft prior to biomechanical testing. With respect to lumbar vertebrae (LV), LV2 are processed for BMD (pQCT may also be performed); LV3 are prepared for undecalcified bone histology; and LV4 are processed for mechanical testing.


Assays - Hair Growth: *In Vivo* Assay of Effects of Compounds on Hair Follicles Proliferation and Hair Growth

The assay described above to assess the effect of compounds on calvarial bone growth can also be used to assess the ability of compounds to stimulate hair growth. The test compound or appropriate control vehicle is applied to the upper and lower back of male ICR Swiss white mice either topically or by subcutaneous injection. The vehicle is selected as appropriate for the compound to be tested and for the route of administration. Optionally, the hair in the test area may be removed prior to administration. After a suitable interval, typically 7 days, the mice are anesthetized and a biopsy of the dorsal treatment area is taken using a 6 mm dermal punch. The specimens are fixed in 10% buffered formalin and imbedded in paraffin wax, and sectioned and stained to observe hair follicles. In addition, photography can be used to observe and record hair growth; typically such growth is observed after 14-18 days. After a suitable interval, typically 21 days, the animals may be euthenized and the hair analyzed for fiber analysis and the tissue from the treatment area analyzed for quantitation of hair follicles.

Nature of the Compounds Useful in the Invention

The compounds useful in the methods and compositions of the invention are inhibitors of proteasomal activity, of the transcription factor NF- κ B, preferably both. Known inhibitors of these activities can be ascertained from the literature or compounds can be tested for these activities using assays known in the art. In addition, inhibitors which lower the level of effective expression of the nucleotide sequence encoding the enzymes that have proteasomal activity or of the nucleotide sequence encoding NF- κ B can be assessed and used in the invention methods.

The compounds thus identified, which are used according to the method of the invention as it relates to treating bone defects, however, preferably do not include compounds that inhibit the isoprenoid pathway, such as the statins. A description of these excluded compounds can be found in WO98/25460 and in U.S. Serial No. 09/096,631, both cited above and incorporated herein by reference. For convenience, the isoprenoid pathway referred to is set forth herein in Figure 2. One class of compounds which are inhibitors are the statins which have the formula



wherein X in each of formulas (1) and (2) represents a substituted or unsubstituted alkylene, alkenylene, or alkynylene linker of 2-6C;

Y represents one or more carbocyclic or heterocyclic rings wherein, when Y comprises two or more rings, said rings may be fused; and

R' represents a cation, H or a substituted or unsubstituted alkyl group of 1-6C; and the dotted lines represent optional π -bonds.

These compounds may, however, be used in the method of the invention as it relates to the stimulation of hair growth.

Compounds known to be proteasome or NF- κ B inhibitors include:

Proteasome Inhibitors	
PSI	N-carbobenzoyl-Ile-Glu-(OIBu)-Ala-Leu-CHO
MG-132	N-carbobenzoyl-Leu-Leu-Leu-CHO
MG-115	N-carbobenzoyl-Leu-Leu-Nva-CHO
MG-101 or Calpain Inh I	N-Acetyl-Leu-Leu-norLeu-CHO
ALLM	N-Acetyl-Leu-Leu-Met-CHO
	N-carbobenzoyl-Gly-Pro-Phe-Leu-CHO
	N-carbobenzoyl-Gly-Pro-Ala-Phe-CHO
	N-carbobenzoyl-Leu-Leu-Phe-CHO
	N-carbobenzoyl-Leu-Ala-Leu-CHO
Gilotoxin	
SN50	NLS of NF- κ B MW 2781
Bay 11-7082	
Capsaicin	
PDTC	

See, for example, Vinitsky, A. *et al. J Biol Chem* (1994) 269:29860-29866;

Figueiredo-Pereira, M.E. *et al. J Neurochem* (1994) 63:1578-1581; Wojcik, C. *et al. Eur*

5 *J Cell Biol* (1996) 71:311-318.

In the foregoing list, lactacystin is known to be an irreversible inhibitor of proteasome activity. It binds to the β catalytic subunit and is a specific inhibitor of the 20S proteasome. It also irreversibly inhibits NF- κ B.

10 SN50 is the NLS (nuclear localization sequence) of p50 plus the hydrophobic region of K-FGF. It inhibits the translocation of the NF- κ B active complex to the nucleus.

Certain peptidyl epoxy ketones such as EST are irreversible inhibitors of the proteasomes. MG-132 shows activity against the chymotryptic activity of the 20S protein

without affecting its ATPase or isopeptidase activity and reversibly inhibits NF- κ B activity. MG-115 and MG-341 show similar activities to MG-132. Various other inhibitors of NF- κ B are less active in the ABA assay. These include capsaicin, curcumin, and resveratrol. Other compounds known to inhibit NF- κ B are gliotoxin and PDTTC (1-pyrrrolidine carboxylic acid). Various other compounds such as BAY-11-7082 and BAY-11-7083 as well as calyculin-A inhibit phosphorylation of NF- κ B. Calpain inhibitor inhibits calpain 1 and the proteasome; other compounds such as olomoucine and roscovitine inhibit cdk2 and/or cdk5.

An additional compound shown to be a proteasome inhibitor is pentoxifylline (PTX). Combaret, L. et al. Mol Biol Rep (1999) 26:95-101. It is active in the *in vitro* calvarial assay described above.


As set forth above, in preferred embodiments of the methods of the invention, the identified compounds used in treatment of bone disorders are other than statins and other compounds that inhibit the isoprenoid pathway, typically as shown in Figure 1. In other preferred embodiments, also excluded from use in the methods of treatment of bone disorders of the present invention, are compounds described in PCT applications WO98/17267, WO97/15308, and WO97/48694 cited and incorporated herein by reference hereinabove. However, the use of these compounds in the method to stimulate hair growth according to the invention is not excluded.

The following examples are intended to illustrate but not to limit the invention.

Example 1

High Throughput Screening

Thousands of compounds have been tested in the assay system set forth in U.S. Serial No. 08/458,434, filed 2 June 1995, and incorporated herein by reference. Representative compounds of the invention gave positive responses, while the majority of (unrelated) compounds are inactive. In this screen, the standard positive control was the compound 59-0008 (also denoted "OS8"), which is of the formula:



In more detail, the 2T3-BMP-2-LUC cells, a stably transformed osteoblast cell line described in Ghosh-Choudhury *et al. Endocrinology* (1996) 137:331-39, referenced above, was employed. The cells were cultured using α -MEM, 10% FCS with 1% penicillin/streptomycin and 1% glutamine ("plating medium"), and were split 1:5 once per week. For the assay, the cells were resuspended in a plating medium containing 4% FCS, plated in microtiter plates at a concentration of 5×10^3 cells (in 50 μ l/well), and incubated for 24 hours at 37°C in 5% CO₂. To initiate the assay, 50 μ l of the test compound or the control in DMSO was added at 2X concentration to each well, so that the final volume was 100 μ l. The final serum concentration was 2% FCS, and the final DMSO concentration was 1%. Compound 59-0008 (10 μ M) was used as a positive control.

The treated cells were incubated for 24 hours at 37°C and 5% CO₂. The medium was then removed, and the cells were rinsed three times with PBS. After removal of excess PBS, 25 μ l of 1X cell culture lysing reagent (Promega #E153A) was added to each well and incubated for at least ten minutes. Optionally, the plates/samples could be frozen at this point. To each well was added 50 μ l of luciferase substrate (Promega #E152A; 10 ml Promega luciferase assay buffer per 7 mg Promega luciferase assay substrate). Luminescence was measured on an automated 96-well luminometer, and was expressed as either picograms of luciferase activity per well or as picograms of luciferase activity per microgram of protein.

In this assay, compound 59-0008 (3-phenylazo-1H-4,1,2-benzothiazine) exhibits a pattern of reactivity which is maximal at a concentration of approximately 3-10 μ M. Accordingly, other tested compounds can be evaluated at various

concentrations, and the results compared to the results obtained for 59-0008 at 10 μ M (which value would be normalized to 100). Alternatively, the reactivity of a compound to be tested can be compared directly to a negative control containing no compound.

The control compound 59-0328, which is simvastatin, gives a good response. The known proteasome inhibitors MG-132 and MG-115 also show high activity; MG-132 is

410 nm in a Beckman dual beam spectrophotometer from the OD410 and corrected for protein concentration.

PSI and MG-132 and control compounds/factors bFGF and BMP-2, and a vehicle control were tested in this assay and the calvaria were analyzed histomorphometrically, as described above. Increase in bone area as a function of concentration; the increase in osteoblasts and the enhancement of alkaline phosphatase activity for PSI were measured.

The data show that PSI is as good as, or better than, BMP-2 and bFGF (two "gold standard" agents for bone growth; see Wozney J. *Molec Reprod Dev* (1992) 32:160-67; WO95/24211) for inducing bone formation.

An additional experiment, penoxifylline (PTX) was tested in the foregoing assay. It exhibited the ability to enhance new bone formation in concentrations as low as 0.1 µm. At a concentration of 10 µm, PTX appeared to enhance the new bone are over control by over 100%; at 100 µm, the increase was approximately three (3) times that of control.

Example 3

In vivo Calvarial Bone Growth Data

PSI and MG-132 were assayed *in vivo* according to the procedure described previously (see "In vivo Assay of Effects of Compounds on Murine Calvarial Bone Growth", *supra*). As a control, simvastatin provided a 1.5 fold increase in the number of osteoblasts.

In one experiment, vehicle control, bFGF and varying doses of PSI were tested in the *in vivo* calvarial bone growth assay. The results are reported as a measurement of total bone area, % increase in area over vehicle control, and % increase in new bone width as shown below.

Compound	Total Bone Area (µm ²)	% Increase* in Bone Area Compared with Control	% Increase* in New Bone Width
Control	0.84 ± 0.03		
0.1 mg/kg/day	0.74 ± 0.02	21.7 ± 3.5	19.9 ± 2.0
1 mg/kg/day	0.83 ± 0.02	35.4 ± 3.4	19.9 ± 4.4
5 mg/kg/day	0.79 ± 0.03	32.1 ± 5.6	
		*p < 0.05	*p < 0.001

25

In addition, histological examination showed confirmation of bone growth both when 5 mg/kg/day of PSI was used and 1 mg/kg/day was used.

effective at lower concentrations. Positive responses are also obtained using lactacystin. However, gliotoxin, olomoucine, roscovitine, SN50, PDTC, and capsaicin do not give promising responses.

Example 2

In vitro Bone Formation

Selected compounds and appropriate controls were assayed *in vitro* (ex vivo) for bone formation activity (described above in "Techniques for Neonatal Mouse Calvaria Assay (*in vitro*)). Histomorphometrical assessments of *ex vivo* calvaria were carried out using an OsteoMetrics bone morphology measurement program, according to the manufacturer's instructions. Measurements were determined using either a 10- or 20-fold objective with a standard point counting eyepiece graticule.

New bone formation was determined (using a 10X objective) by measuring the new bone area formed in one field in 3 representative sections of each bone (4 bones per group). Each measurement was carried out 1/4 field distance from the end of the suture. Both total bone and old bone area were measured. Data were expressed as new bone area in µm².

The results in Example 1 were somewhat imperfectly correlated with the results in this assay. The control compound, simvastatin showed new bone formation in this assay as did MG-132 and lactacystin. MG-115 also showed positive results although less dramatic than those of simvastatin. However, gliotoxin, which appeared negative in the ABA assay of Example 1 did demonstrate the ability to stimulate bone growth. The remaining compounds, olomoucine, roscovitine, SN50, PDTC and capsaicin appeared negative in this assay.

Osteoblast numbers are determined by point counting. The number of osteoblast cells lining the bone surface on both sides of the bone are counted in one field using a 20X objective. Data are expressed as osteoblast numbers/mm of bone surface.

Alkaline phosphatase activity is measured in the conditioned media of the murine organ cultures, using the method described by Majeska, R.J. *et al. Exp Cell Res* (1978) 111:465-465. Conditioned media are incubated at 37°C for 20 minutes with phosphatase substrate 104 (Sigma) and the reaction stopped with 2 ml of 0.1 M NaOH. Alkaline phosphatase activity is calculated by measuring cleaved substrate at an optical density of






30








Example 4**Summary of Effects on Bone Formation**

The table below summarizes the results obtained for compounds tested in the various assays set forth above. It is seen that compounds that are proteasome inhibitors also enhance bone formation. In the compounds tested in this table, however, compounds which are known to be inhibitors only of NF- κ B but which fail to inhibit proteasomal activity, do not enhance luciferase activity (indicative of BMP-2 promoter activity) in the high through-put assay, nor do they enhance bone formation in the calvarial assay *in vitro*, to as great an extent as do proteasome inhibitors.

10

Compounds useful in the invention include:

Compound	Structure	Luciferase Activity (ED ₅₀ - μ M)	Bone Formation (ED ₅₀ - μ M)	Proteasome Activity (ED ₅₀ - μ M)
Simvastatin		\cap 0.2	\cap 0.2	\cap
Lactacystin		\cap 1	\cap 1	\cap 1.5
PS1	Z-Ile-Glu(OiBu)-Ala-Leu-CHO	\cap 0.05	\cap 0.03	\cap 0.035
MG132		\cap 0.25	\cap 0.5	\cap 0.3
MG262		\cap 0.1	\cap 0.1	\cap 0.07
MG115		\cap 2	\cap 1	\cap 1

Compound	Structure	Luciferase Activity (ED ₅₀ - μ M)	Bone Formation (ED ₅₀ - μ M)	Proteasome Activity (ED ₅₀ - μ M)
ALLN	 N-Acetyl-Leu-Leu-Nle-CHO	\cap 10	\cap	\cap 1.5
Cyclosporin A		\cap	\cap 10	\cap 1.0
Gliotoxin		\cap	\cap 10	\cap
SN50	NLS of NF- κ B MW 2781	\cap	\cap	\cap
ALLM	N-Acetyl-Leu-Leu-Meth-CHO	\cap	\cap	\cap 4
PPM-18		\cap	\cap	\cap
Bay 11-7082		\cap	\cap	\cap
Capsaicin		\cap	\cap	\cap 30
PDTC		\cap	\cap	\cap

Example 5Effect of PSI and Other Proteasome Inhibitors on Hair Follicle Production

The *in vivo* bone calvarial growth assay of Example 3 was modified to observe the number of hair follicles in treated mice. In initial observations, PSI (5mg/kg/day) was injected three times a day for 5 days over the calvaria of Swiss ICR mice as described above. Sixteen days later the mice were sacrificed. Histology of the calvaria revealed a strikingly large increase in the number of hair follicles in those mice treated with PSI versus control mice. In addition to PSI, MG132 (10mg/kg), MG115 (10mg/kg) and lactacystin administered in the same way also stimulated an increase in the number of hair follicles.

10

Example 6Stimulation of Hair Growth

Male Swiss ICR mice were first treated to remove hair from the scalp and dorsal regions as follows. Paraffin wax was liquefied by heating to 55°C and the liquefied wax then applied by brush to the scalp and/or back (under anesthesia). The wax was allowed to solidify and then removed. The day following hair stripping, PSI (1 mg/kg/day) was injected subcutaneously three times a day for five days into the scalp and dorsal region. On day 7 a dermal punch biopsy was taken; histology revealed a large increase in the number of hair follicles in mice administered PSI versus control mice. By day 18, it was observable that the treated mice had a hair growth rate greater than that of the mice in the control group.

The mice were sacrificed on day 21 and histology was performed on the dermis of the scalp and of the dorsal region. In the treated mice, mature hair follicles in numbers much greater than in controls had migrated to the lower region of the dermis. Upon closer examination, it was observed that mice that had received only vehicle had quiescent hair follicles. When treated with PSI such follicles were stimulated to differentiate into mature hair follicles and to migrate to the lower region of the dermis.

Example 7Topical Administration

PSI was prepared as a topical formulation, where the vehicle was 50% propylene glycol, 30% ethanol, 20% deionized water, at 0.1% concentration of PSI. The solution

30

was applied 3 times a day for 5 days. The mice in a treated group were observed as compared to controls similarly treated with vehicle alone. The results at day 16 showed stimulation of hair growth relative to the controls.

In addition to stimulating hair growth, PSI was able to thicken both the hair and the hair shaft. PSI increases hair count when the follicle area is greater than 0.01 mm². When the protocol above was repeated using a 0.5% solution of PSI in groups containing 5 mice each, the number of hairs per 0.8 mm² was 60 in the treated mice versus about 10 in the control group. The percentage of follicle area in a region of about 0.8 mm² was about 30% as an average in the treated group as compared to 15% as an average in the control group.

10

Example 8Dose Requirements

In order to determine the minimal effective dosage of PSI, when used topically, a dose response curve for PSI was prepared. All experiments were performed according to current good laboratory practice regulations (21CFR58). The mice were divided into 7 groups, 10 mice each, wherein one group was control treated only with vehicle and groups 1-6 with a series of increasing concentrations of PSI in a vehicle comprising 50% propylene glycol, 30% ethanol, 20% deionized water. The concentrations were 0.006%, 0.012%, 0.025%, 0.05%, 0.1% and 0.5%.

20

The mice were anesthetized (50 µl Mouse Cocktail containing 3 ml ketamine, 2 ml small animal rompun, 5 ml NaCl), identified by ear punch code, weighed and the hair on the dorsal side removed by waxing as described in Example 6. After waxing, the animals were photographed. On the following day (day 1), 100 µl of PSI at the above concentrations in vehicle was brushed onto the area of removed hair. A similar application of PSI solution was performed daily for an additional 4 days.

25

On Day 7 mice were anesthetized and a biopsy of the dorsal treatment area taken using a 6 mm dermal punch; the specimens were fixed in 10% buffered formalin and embedded in paraffin wax. Sections were cut using a standard microtome.

30

Mice were monitored daily for signs of hair growth, and any hair growth was recorded by photography. On day 21 animals were euthanized (75mg/kg body weight phenobarbital, IP injection), 2 cm hair samples were taken for optical based fiber analysis, and the remaining dorsal treatment area was fixed in 10% buffered formalin for further

histological analysis. Analysis included quantification of hair thickness and quantification of mature hair follicles. Results were expressed as the mean \pm standard error of the mean. Data were analyzed by repeated measures of analysis of variance followed by the Tukey-Kramer post test. P values of <0.05 were considered significant.

The results indicate that the minimal effective dose of PSI is 0.5% applied 1 time a day for 4 days; additional experiments showed that 0.1% of PSI applied topically 3 times a day for 5 days was also effective.

Gross observation of mice receiving an effective dose indicated an enhanced rate of hair growth, a thickening of hair diameter, increase in sheath diameter, and differentiation of quiescent hair follicles into more mature forms.

The contents of all documents cited above are expressly incorporated herein to the extent required to understand the invention.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

Claims

1. A method to enhance bone formation or to treat pathological dental conditions or to treat degenerative joint conditions in a vertebrate animal which method comprises administering to a vertebrate subject in need of such treatment an effective amount of a compound that inhibits the activity of NF- κ B or that inhibits proteasomal activity or that inhibits production of proteasome proteins.

2. The method of claim 1 wherein said compound inhibits proteasomal activity or that inhibits production of proteasomal proteins.

3. The method of claim 1 wherein said compound does not inhibit the isoprenoid pathway.

4. The method of claim 1 wherein said compound is lactacystin, a peptidyl aldehyde, or PTX.

5. The method of claim 1 wherein said subject is characterized by a condition selected from the group consisting of osteoporosis, bone fracture or deficiency, primary or secondary hyperparathyroidism, periodontal disease or defect, metastatic bone disease, osteolytic bone disease, post-plastic surgery, post-prosthetic joint surgery, and post-dental implantation.

6. The method of claim 1 which further comprises administering to said subject one or more agents that promote bone growth or that inhibit bone resorption.

7. The method of claim 6 wherein said agents are selected from the group consisting of bone morphogenetic factors, anti-resorptive agents, osteogenic factors, cartilage-derived morphogenetic proteins, growth hormones, estrogens, bisphosphonates, statins and differentiating factors.

8. A method to treat a mammalian subject for a condition benefited by stimulating hair growth which method comprises administering to said mammalian

subject in need of such treatment an effective amount of a compound that inhibits the activity of NF- κ B or that inhibits proteasomal activity or that inhibits production of these proteins.

9. The method of claim 8 wherein said compound inhibits proteasomal activity or that inhibits production of proteasome proteins.

10. The method of claim 9 wherein said compound is lactacystin or a peptidyl aldehyde.

11. A pharmaceutical composition for treating bone disorders, dental pathological conditions or degenerative joint conditions which composition comprises a compound that inhibits the activity of NF- κ B or that inhibits proteasomal activity or that inhibits production of these proteins.

12. The pharmaceutical composition of claim 11 wherein said compound inhibits proteasomal activity or that inhibits production of proteasomal proteins.

13. The pharmaceutical composition of claim 11 wherein said compound does not inhibit the isoprenoid pathway.

14. The pharmaceutical composition of claim 11 wherein said compound is lactacystin, a peptidyl aldehyde, or PTX.

15. The pharmaceutical composition of claim 11 wherein said compound does not inhibit the isoprenoid pathway.

16. A pharmaceutical composition for treating for a condition benefited by stimulating hair growth which composition comprises a compound that inhibits the activity of NF- κ B or that inhibits proteasomal activity or that inhibits production of these proteins.

17. The pharmaceutical composition of claim 13 wherein said compound is lactacystin or a peptidyl aldehyde.

18. A method to identify a compound which enhances bone growth or stimulates hair growth which method comprises subjecting said compound to an assay for determining its ability to inhibit NF- κ B activity, whereby a compound which inhibits the activity of NF- κ B is identified as a compound which enhances bone growth; or

subjecting said compound to an assay for determining its ability to inhibit the production of NF- κ B, whereby a compound which inhibits the production of NF- κ B is identified as a compound which enhances bone growth; or

subjecting a candidate compound to an assay to assess its ability to inhibit proteasomal activity, whereby a compound which inhibits proteasomal activity is identified as a compound that enhances bone growth; or

subjecting a candidate compound to an assay to assess its ability to inhibit the

production of enzymes with proteasomal activity, whereby a compound which inhibits the production of enzymes with proteasomal activity is identified as a compound that enhances bone growth.




Figure 1

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.